arXiv:1010.3181v2 [hep-ph] 28 Jul 2011

The Effects of a Family Non-universal Z' Boson in the $\bar{B}_s \to \pi K$, πK^*, ρK Decays and $B_d - \bar{B}_d$ Mixing

Qin Chang[a,b]*, Ya-Dong Yang[b,c]

[a]Department of Physics, Henan Normal University, Xinxiang, Henan 453007, P. R. China

[b]Institute of Particle Physics, Huazhong Normal University, Wuhan, Hubei 430079, P. R. China

[c]Key Laboratory of Quark & Lepton Physics, Ministry of Education, P.R. China

Abstract

We revisit $\bar{B}_s \to \pi K$, πK^* and ρK decays within QCDF formalism and examine the possible effects of a family non-universal Z' boson in these decays. In our evaluations, the strong constraints from $B_d - \bar{B}_d$ mixing on the strength of the left-handed flavor-changing $d - b - Z'$ coupling B_{db}^L ($\sim 0.2 \times 10^{-3}$) is also included. Numerically, we find that a new weak phase $\phi_d^L \sim -50^\circ$ involved in B_{db}^L, the negative combination of the $u - u - Z'$ and $d - d - Z'$ couplings P_{ud}^R and/or D_{ud}^L with larger absolute value are crucial to improve the agreement of $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ between the SM prediction and the experimental measurement. Generally, $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ could be reduced by about 18% at most by Z' contributions. Moreover, combining with the recent updated measurement on β_s by CDF and D0 collaborations, we also find that Z' effects in $\bar{B}_s \to \pi^- K^+$ decays are very important for examining whether the CKM-like hierarchy persists in a family non-universal Z' coupling matrix.

PACS Numbers: 13.25.Hw, 12.60.Cn, 12.15.Mm, 11.30.Hv.

---

*Corresponding author. changqin@htu.cn

1 Introduction

The fruitful running of BABAR, Belle, CDF and D0 in the past years provides a very fertile testing ground for the Standard Model (SM) picture of flavor physics and CP violations. Although most of the measurements are in perfect agreement with the SM predictions, there still exist some unexplained mismatches, such as the so-called "πK, $\pi\pi$ puzzles", anomalous $\bar{B}_s - B_s$ mixing phase, large transversal polarization in $B \to \phi K^*$, mismatching forward-backward asymmetry in $B \to K^* \mu^+ \mu^-$ and so on. Recently, some interesting measurements of B_s decays have been reported by CDF collaboration at Fermilab Tevatron, which have attracted many attentions. Especially, CDF collaboration has made the first measurement of $\bar{B}_s \to \pi^- K^+$ decay [1, 2]

$$\begin{cases} \mathcal{B}(\bar{B}_s \to \pi^- K^+) = (5.0 \pm 0.7 \pm 0.8) \times 10^{-6}\,, \\ A_{cp}^{dir}(\bar{B}_s \to \pi^- K^+) = (39 \pm 15 \pm 8)\%\,. \end{cases} \quad \text{CDF Collaboration} \tag{1}$$

The recent theoretical predictions for these two quantities based on the QCD factorization approach (QCDF)[3], the perturbative QCD approach (pQCD)[4] and the soft-collinear effective theory (SCET) [5] read

$$\begin{cases} \mathcal{B}(\bar{B}_s \to \pi^- K^+)_{QCDF} = 8.3 \times 10^{-6}\,, \\ A_{cp}^{dir}(\bar{B}_s \to \pi^- K^+)_{QCDF} = 10.9\%\;; \end{cases} \quad \text{QCDF Scenario S4 [6]} \tag{2}$$

$$\begin{cases} \mathcal{B}(\bar{B}_s \to \pi^- K^+)_{pQCD} = (7.6^{+3.2}_{-2.3} \pm 0.7 \pm 0.5) \times 10^{-6}\,, \\ A_{cp}^{dir}(\bar{B}_s \to \pi^- K^+)_{pQCD} = (24.1^{+3.9+3.3+2.3}_{-3.6-3.0-1.2})\%\,; \end{cases} \quad \text{pQCD [7]} \tag{3}$$

$$\begin{cases} \mathcal{B}(\bar{B}_s \to \pi^- K^+)_{SCET} = (4.9 \pm 1.2 \pm 1.3 \pm 0.3) \times 10^{-6}\,, \\ A_{cp}^{dir}(\bar{B}_s \to \pi^- K^+)_{SCET} = (20 \pm 17 \pm 19 \pm 5)\%. \end{cases} \quad \text{SCET [8]} \tag{4}$$

One may note that QCDF (Scenario S4) result for $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ is significantly larger than CDF measurement Eq. (1). The pQCD prediction agree with the data in Eq. (1) within its large theoretical uncertainty, even though its central value also much larger than the measurement Eq. (1). In SCET, the contributions involving internal charm quark loops are claimed to be non-perturbative and only can be determined by fitting to the data. A recent fit analysis of charmless B decays using SCET gives the result Eq. (4), which agrees with CDF data. All of the theoretical predictions for $A_{cp}^{dir}(\bar{B}_s \to \pi^- K^+)$ agree with CDF data with their large theoretical uncertainties.

For this possible branching ratio mismatch, many efforts have been done within both SM and possible New Physics (NP) scenarios, for example Refs. [9, 10, 11, 12]. A recent detailed analysis in Ref. [9] indicates that $SU(3)$ and factorization only remain approximately valid if the branching ratio for $\bar{B}_s \to \pi^- K^+$ exceeds its current value by at least 42%, or if a parameter ξ describing ratios of form factors and decay constants is shifted from its nominal value by more than twice its estimated error. Such analysis also implies that, if $\mathcal{B}_{Exp}(\bar{B}_s \to \pi^- K^+)$ and ξ persist in the future, large $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ mismatch is possibly induced by NP effects.

A new family non-universal Z' boson could be naturally derived in certain string constructions [13], E_6 models [14] and so on. Searching for such an extra Z' boson is an important mission in the experimental programs of Tevatron [15] and LHC [16]. Within a model-independent formalism for a family non-universal Z' model presented by Ref. [17], $b \to s$ transition involving $b - s - Z'$ coupling has been widely studied. It is interesting that such Z' boson behavior is helpful to resolve many puzzles in $b \to s$ transition, such as "πK puzzle" [18, 19] and anomalous $\bar{B}_s - B_s$ mixing phase [20, 21]. So, it is worth evaluating the effects of a non-universal Z' boson on $b \to d$ transition, especially the measured $\bar{B}_s \to \pi^- K^+$. Moreover, with the constraints from $\bar{B}_q - B_q$ mixing, $B \to \pi K$, πK^*, ρK and $X_s \mu^+ \mu^-$ decays, a CKM-like hierarchy persists in Z' coupling matrix (at least between $b - s - Z'$ and $b - d - Z'$ couplings) [21]. Naturally, it is important to examine whether such hierarchy is persisted after the constraints from $\bar{B}_s \to \pi^- K^+$ are included.

In addition, the correlated decays $\bar{B}_s \to \pi K^*$ and ρK, which also involves $b \to d\bar{q}q$ $(q = u, d)$ transition at quark level, will be measured at the running LHC. Such decays may perform an important role in testing SM and searching for NP signal. Especially, due to large theoretical uncertainty cancellation, the ratio $\mathcal{B}(\bar{B}_s \to \rho^- K^+)/\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ is much more suitable for testing SM and probing NP [10]. Therefore, we also evaluate $\bar{B}_s \to \pi K^*$ and ρK decays within both SM and a family non-universal Z' model.

Our paper is organized as follows. In Section 2, we revisit $\bar{B}_s \to \pi K$, πK^* and ρK decays in the SM within the QCDF formalism; we mainly evaluate the effects of annihilation contributions which involve endpoint divergency. In Section 3, we perform a fitting on $b - d - Z'$ couplings with the constraints from $\bar{B}_d - B_d$ mixing and $\bar{B}_s \to \pi^- K^+$ decay; the NP effects on the other decays are also evaluated. Section 4 is our conclusions. Theoretical input parameters are summarized in the Appendix.

2 Revisiting $\bar{B}_s \to \pi K$, πK^* and ρK Decays within QCDF Formalism

In the SM, the effective weak Hamiltonian responsible for $b \to d$ transitions is given as [24]

$$\begin{aligned}\mathcal{H}_{\rm eff} &= \frac{G_F}{\sqrt{2}}\biggl[V_{ub}V_{ud}^*\,(C_1O_1^u + C_2O_2^u) + V_{cb}V_{cd}^*\,(C_1O_1^c + C_2O_2^c) - V_{tb}V_{td}^*\,\bigl(\sum_{i=3}^{10} C_iO_i \\ &\quad + C_{7\gamma}O_{7\gamma} + C_{8g}O_{8g}\bigr)\biggr] + \text{h.c.},\end{aligned} \tag{5}$$

where $V_{qb}V_{qd}^*$ ($q = u, c$ and t) are products of the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements [25], C_i the Wilson coefficients, and O_i the relevant four-quark operators.

In recent years, QCDF has been employed extensively to study the B meson non-leptonic decays. For example, all of the decay modes considered here have been studied comprehensively within the SM in Refs. [6, 11]. The relevant decay amplitudes for $\bar{B}_s \to \pi K$, πK^* and ρK decays within the QCDF formalism are

$$\begin{aligned}\mathcal{A}_{\bar{B}_s\to\pi^-K^+} &= A_{K\pi}\left[\delta_{pu}\alpha_1 + \alpha_4^p + \alpha_{4,\rm EW}^p + \beta_3^p - \frac{1}{2}\beta_{3,\rm EW}^p\right], \\ \sqrt{2}\mathcal{A}_{\bar{B}_s\to\pi^0K^0} &= A_{K\pi}\left[\delta_{pu}\alpha_2 - \alpha_4^p + \frac{3}{2}\alpha_{3,\rm EW}^p + \frac{1}{2}\alpha_{4,\rm EW}^p - \beta_3^p + \frac{1}{2}\beta_{3,\rm EW}^p\right].\end{aligned} \tag{6}$$

where the explicit expressions for the coefficients $\alpha_i^p \equiv \alpha_i^p(M_1M_2)$ and $\beta_i^p \equiv \beta_i^p(M_1M_2)$ can also be found in Ref. [6]. The expressions for the $\bar{B}_s \to \pi K^*$ and ρK amplitudes are obtained by setting $(\pi K) \to (\pi K^*)$ and $(\pi K) \to (\rho K)$, respectively.

It is noted that the QCDF framework contains estimates of some power-suppressed but numerically important contributions, such as the annihilation corrections. Unfortunately, in a collinear factorization approach, the calculation of annihilation amplitude always suffers from end-point divergence. In the pQCD approach, such divergence is regulated by the parton transverse momentum k_T at the expense of modeling additional k_T dependence of meson distribution functions [4], and a large strong phase is found. In Refs. [8, 26], annihilation diagram is studied with SCET and parameterized by a complex amplitude. At present, the dynamical origin of these corrections still remains a theoretical challenge. Within the QCDF framework, in a model-dependent way, there are two schemes to regulate the endpoint divergence until now. Scheme I: In Ref. [6], to probe their possible effects conservatively, the endpoint divergent integrals are treated as signs of infrared sensitive contribution and phenomenologically parameterized by

$$\int_0^1 \frac{dx}{x} \to X_{A,H} = (1+\rho_{A,H}e^{i\phi_{A,H}}) \ln\frac{m_B}{\Lambda_h}, \qquad \int_0^1 dy \frac{\ln y}{y} \to -\frac{1}{2}(X_{A,H})^2 \tag{7}$$

with $\rho_{A,H} \leq 1$ and $\phi_{A,H}$ unrestricted. The different scenarios corresponding to different choices of $\rho_{A,H}$ and $\phi_{A,H}$ have been thoroughly discussed in Ref. [6]. Scheme II: As an alternative scheme to the first one, one could quote an infrared finite gluon propagator to regulate the endpoint divergent integrals, which have been thoroughly studied in Ref. [28].



(a) (b)

Figure 1: The dependence of $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ on $\phi_{A,H}$ with $\rho_{A,H} = 1$ (colored real line), 2(colored dashed line). The black horizontal lines are the experimental data with the solid line being the central value and the dashed ones the error-bars (1σ).

In scheme I, with the central value of the input parameters listed in the Appendix, Fig. 1 shows the dependence of $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ on $\phi_{A(H)}$ with $\rho_{A(H)} = 1$, 2 and $\rho_{H(A)} = 0$. As Fig. 1(a) shows, assuming $\phi_A \neq \phi_H$, the minimal $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ appears at $\phi_H \simeq 0°$ and $\phi_A \simeq -110°$; Assuming $\phi_A = \phi_H$, it appears at $\phi_H = \phi_A \simeq -100°$, as shown in Fig. 1 (b). With the above $\phi_{H,A}$ values, $\rho_{A,H} = 1$ and the central values of the other inputs, corresponding above two assumptions for $\phi_{A,H}$, we get $\mathcal{B}(\bar{B}_s \to \pi^- K^+) = 9.0 \times 10^{-6}$, 9.3×10^{-6} respectively, which are also significantly larger than CDF data $(5.0 \pm 1.1) \times 10^{-6}$. Moreover, corresponding to above two assumptions, we find the direct CP asymmetry for $\bar{B}_s \to \pi^- K^+$ is 3%, 6% respectively, which are also lower than CDF measurement $(39 \pm 17)\%$ at 2σ level. So, following

Table 1: The numerical results for $\mathcal{B}(\bar{B}_s \to \pi K, \pi K^*, \rho K)(\times 10^{-6})$.

	Exp.	SM		SM+Z'	
		Scheme I	Scheme II	S1	S2
$\mathcal{B}(\bar{B}_s \to \pi^- K^+)$	5.0 ± 1.1	$9.5^{+3.8}_{-2.6}$	$10.2^{+3.7}_{-2.5}$	$8.6^{+3.8}_{-2.8}$	$8.9^{+3.9}_{-2.6}$
$\mathcal{B}(\bar{B}_s \to \pi^0 K^0)$	—	$0.53^{+0.35}_{-0.12}$	$0.39^{+0.23}_{-0.08}$	$0.50^{+0.40}_{-0.17}$	$0.52^{+0.41}_{-0.20}$
$\mathcal{B}(\bar{B}_s \to \pi^- K^{*+})$	—	$13.3^{+4.5}_{-3.4}$	$14.1^{+4.0}_{-3.4}$	$14.0^{+4.6}_{-3.9}$	$13.7^{+5.2}_{-3.6}$
$\mathcal{B}(\bar{B}_s \to \pi^0 K^{*0})$	—	$0.38^{+0.23}_{-0.15}$	$0.28^{+0.16}_{-0.06}$	$0.21^{+0.27}_{-0.10}$	$0.26^{+0.16}_{-0.15}$
$\mathcal{B}(\bar{B}_s \to \rho^- K^+)$	—	$22.5^{+9.2}_{-6.5}$	$24.5^{+9.1}_{-6.4}$	$22.2^{+10.2}_{-6.6}$	$22.2^{+9.2}_{-6.4}$
$\mathcal{B}(\bar{B}_s \to \rho^0 K^0)$	—	$0.79^{+0.42}_{-0.35}$	$0.27^{+0.09}_{-0.06}$	$0.59^{+0.66}_{-0.37}$	$0.64^{+0.58}_{-0.37}$

Table 2: The numerical results for $A_{cp}^{dir}(\bar{B}_s \to \pi K, \pi K^*, \rho K)(\times 10^{-2})$.

	Exp.	SM		SM+Z'	
		Scheme I	Scheme II	S1	S2
$A_{cp}^{dir}(\bar{B}_s \to \pi^- K^+)$	39 ± 17	$13.6^{+6.4}_{-4.6}$	$22.4^{+7.6}_{-5.7}$	$14.2^{+6.8}_{-5.7}$	$14.0^{+7.6}_{-5.2}$
$A_{cp}^{dir}(\bar{B}_s \to \pi^0 K^0)$	—	$19.7^{+10.6}_{-15.5}$	$10.4^{+7.9}_{-7.8}$	$25.7^{+15.5}_{-16.0}$	$23.8^{+16.3}_{-14.3}$
$A_{cp}^{dir}(\bar{B}_s \to \pi^- K^{*+})$	—	$-17.1^{+5.7}_{-4.8}$	$-23.9^{+7.7}_{-6.5}$	$-16.5^{+5.7}_{-4.3}$	$-16.7^{+5.7}_{-5.1}$
$A_{cp}^{dir}(\bar{B}_s \to \pi^0 K^{*0})$	—	$-15.0^{+10.6}_{-26.1}$	$-16.2^{+7.1}_{-13.8}$	-50^{+42}_{-34}	-34.9^{+30}_{-28}
$A_{cp}^{dir}(\bar{B}_s \to \rho^- K^+)$	—	$9.6^{+2.7}_{-5.4}$	$18.3^{+4.6}_{-4.8}$	$9.6^{+1.8}_{-5.4}$	$9.6^{+1.7}_{-5.2}$
$A_{cp}^{dir}(\bar{B}_s \to \rho^0 K^0)$	—	$3.0^{+18.8}_{-8.5}$	$-19.9^{+16.3}_{-10.9}$	23.7^{+44}_{-23}	$16.0^{+38.2}_{-18.8}$

a reasonable choice presented by Cheng and Chua [11], in our following calculations, we take $\rho_{A,H}(PP, PV, VP) = 1$ and $\phi_{A,H} = -55°(PP)$, $-30°(PV)$, $-65°(VP)$. Such choice of the values of parameters $\rho_{A,H}$ and $\phi_{A,H}$ are similar to the so-called favored scenario "scenario S4" for $B_{u,d}$ decays presented by Beneke and Neubert [6], except that some additional moderate values of input parameters (such as $m_s = 80\,MeV$ and $\alpha_2^{\pi} = 0.3$) are also used in "scenario S4" [6]. For estimating theoretical uncertainties, we shall assign an error of ± 0.1 to $\rho_{A,H}$ and $\pm 5°$ to $\phi_{A,H}$. With such $\rho_{A,H}$, $\phi_{A,H}$ values and the other inputs listed in the Appendix, we present the QCDF predictions for $\bar{B}_s \to \pi K$, πK^* and ρK decays in the third column of Table 1, 2 and 3.

Table 3: The numerical results for $A_{CP}^{mix}(\bar{B}_s \to \pi^0 K_s)$ and $A_{CP}^{mix}(\bar{B}_s \to \rho^0 K^0)(\times 10^{-2})$.

	Exp.	SM		SM+Z'	
		Scheme I	Scheme II	S1	S2
$A_{CP}^{mix}(\bar{B}_s \to \pi^0 K_s)$	—	-83^{+19}_{-13}	$-96^{+2.9}_{-1.8}$	-89^{+23}_{-8}	-88^{+20}_{-93}
$A_{CP}^{mix}(\bar{B}_s \to \rho^0 K^0)$	—	-22^{+35}_{-28}	-78^{+12}_{-7}	-56^{+59}_{-31}	-46^{+48}_{-41}

In scheme II, we adopt the infrared finite gluon propagator derived by Cornwall [27], which involves a new parameter gluon mass scale m_g, to regulate the endpoint divergency. In Ref. [28], we present our suggestion $m_g \sim 500$ MeV, which is a reasonable choice so that most of the observables for $B \to \pi K$, πK^* and ρK decays are in good agreement with the experimental data. Furthermore, compared to the available data for $B^0 \to K^+ K^-$,$D_s^{(*)} K$, $B_s \to \pi\pi$ and so on, the most recent evaluation [29] also presents that the gluon mass scale is close to 500 MeV. So, in this paper, we shall take $m_g = 500$ MeV. With such a value, the predictions for observables of $\bar{B}_s \to \pi K$, πK^* and ρK decays are listed in the fourth column in Table 1, 2 and 3. We find most of the predictions in scheme I and scheme II are consistent with each other except for some observables in $\bar{B}_s \to \rho^0 K^0$, $\pi^0 K^{*0}$ decays. Because the cancellations among α_2 and $-\alpha_4^p + \frac{3}{2}\alpha_{3,\rm EW}^p + \frac{1}{2}\alpha_{4,\rm EW}^p$ in Eq. (6) for $\bar{B}_s \to \rho^0 K^0$, $\pi^0 K^{*0}$ decays, the effect of annihilation contributions is significant. Furthermore, in $B \to PV$ decay, the annihilation contributions in scheme II provide a larger imaginary part than the one in scheme I [28]. So, the two schemes present some different predictions for $\bar{B}_s \to \rho^0 K^0$, $\pi^0 K^{*0}$ decays, which will be judged by the upcoming LHC-b and proposed super-B experiments.

Within both schemes I and II, we find that $\mathcal{B}_{SM}(\bar{B}_s \to \pi^- K^+)$ is significantly larger than the measurement $(5.0 \pm 1.1) \times 10^{-6}$. Due to the dominance of the tree contribution α_1 in the amplitude for $\bar{B}_s \to \pi^- K^+$ decay, the contributions from annihilations are tiny in comparison to α_1. So, the large $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ problem could not be resolved by tuning the annihilation contributions.

A recent detailed analysis [9] in a flavor symmetry framework indicate that the mismatch $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ problem persists within the SM if experimental result of $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ would not exceeds its current value $(5.0 \pm 1.1) \times 10^{-6}$ [1, 2] by at least 50% or if the SU(3)-breaking factor defined by

$$\xi \equiv \frac{f_K F_{B^0\pi}(m_K^2)}{f_\pi F_{B_s K}(m_\pi^2)} \frac{m_{B^0}^2 - m_\pi^2}{m_{B_s}^2 - m_K^2} \tag{8}$$

is not shifted from the value $0.97^{+0.09}_{-0.11}$ [9, 22] (corresponding to almost exact $SU(3)$, see Refs. [9, 22] for detail) by more than twice its estimated error. With the central value $F_0^{B_s K} = 0.24$ obtained by lattice and pQCD calculations, recent QCDF analysis [11] gets a good result $\mathcal{B}(\bar{B}_s \to \pi^- K^+) \approx 5.3 \times 10^{-6}$. However, as mentioned in Ref. [11], their result $\xi = 1.24$ is larger than 0.97 at 3σ. Furthermore, $F_0^{B_s K}$ is predicted by lattice QCD computations at small recoil, while unfortunately charmless B decays happen at large recoil. So, if the future refined measurement and theoretical study confirm the large $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ problem, it will be a significant signal of NP. In our following study, we will examine the effects of a family non-universal Z' model for possible solution.

3 The effects of family non-universal Z' model

3.1 Brief review of the family non-universal Z' model

A possible heavy Z' boson is predicted in many extensions of the SM, such as grand unified theories, superstring theories, and theories with large extra dimensions. The simplest way to extend the SM gauge structure is to include a new $U(1)$ gauge group. A family non-universal Z' boson can lead to FCNC processes even at tree level due to the non-diagonal chiral coupling matrix. The formalism of the model has been detailed in Ref. [17]. The relevant studies in the context of B physics have also been extensively performed in Refs. [18, 20, 30, 31].

With the assumption of flavor-diagonal right-handed couplings , the Z' part of the effective Hamiltonian for $b \to d\bar{q}q$ $(q = u, d)$ transitions can be written as [19, 20]

$$\mathcal{H}_{eff}^{Z'} = -\frac{G_F}{\sqrt{2}} V_{td}^* V_{tb} \sum_q (\Delta C_3 O_3^q + \Delta C_5 O_5^q + \Delta C_7 O_7^q + \Delta C_9 O_9^q) + h.c. \, , \tag{9}$$

where $O_i^q (i = 3, 5, 7, 9)$ are the effective operators in the SM, and ΔC_i the modifications to the corresponding SM Wilson coefficients caused by Z' boson. In terms of the model parameters

Table 4: The inputs of the Z' couplings [21].

Solutions	P_{ud}^{L}	P_{ud}^{R}	D_{ud}^{L}	D_{ud}^{R}
S1	2.1 ± 3.3	-0.3 ± 1.3	-0.5 ± 0.4	0.03 ± 0.10
S2	0.7 ± 1.9	-0.1 ± 0.8	-0.3 ± 0.2	0.01 ± 0.05

at the M_W scale, ΔC_i are expressed as

$$\begin{aligned} \Delta C_{3,5} &= -\frac{2}{3V_{td}^{*}V_{tb}}\, B_{db}^{L}\,(B_{uu}^{L,R} + 2B_{dd}^{L,R})\,, \\ \Delta C_{9,7} &= -\frac{4}{3V_{td}^{*}V_{tb}}\, B_{db}^{L}\,(B_{uu}^{L,R} - B_{dd}^{L,R})\,, \end{aligned} \tag{10}$$

where $B_{q'q}^{L,R}$ is the effective chiral Z' coupling matrix element.

Generally, the diagonal elements of the effective coupling matrix $B_{qq}^{L,R}$ are real as a result of the hermiticity of the effective Hamiltonian. However, the off-diagonal one B_{db}^{L} may contain a new weak phase ϕ_d^L. Then, conveniently we can represent ΔC_i as

$$\begin{aligned} \Delta C_{3,5} &= 2\,\frac{|V_{td}^{*}V_{tb}|}{V_{td}^{*}V_{tb}}\,\zeta^{LL,LR}\,e^{i\phi_d^L}\,, \\ \Delta C_{9,7} &= 4\,\frac{|V_{td}^{*}V_{tb}|}{V_{td}^{*}V_{tb}}\,\xi^{LL,LR}\,e^{i\phi_d^L}\,, \end{aligned} \tag{11}$$

where the real NP parameters $\zeta^{LL,LR}$, $\xi^{LL,LR}$ and ϕ_d^L are defined, respectively, as

$$\begin{aligned} \zeta^{LL,LR} &= -\frac{1}{3}\,\Big|\frac{B_{db}^{L}}{V_{td}^{*}V_{tb}}\Big|\,P_{ud}^{L,R}\,, \quad P_{ud}^{L,R} = (B_{uu}^{L,R} + 2B_{dd}^{L,R})\,; \\ \xi^{LL,LR} &= -\frac{1}{3}\,\Big|\frac{B_{db}^{L}}{V_{td}^{*}V_{tb}}\Big|\,D_{ud}^{L,R}\,, \quad D_{ud}^{L,R} = (B_{uu}^{L,R} - B_{dd}^{L,R})\,; \\ \phi_d^L &= \mathrm{Arg}[B_{db}^{L}]\,. \end{aligned} \tag{12}$$

Firstly, we shall specify the values of the Z' input parameters in our calculations. In $b \to s\bar{q}q$ transition, the combinations of $b-s-Z'$, $u-u-Z'$ and $d-d-Z'$ couplings (*i. e.* $\zeta_s^{LL,LR}$ and $\xi_s^{LL,LR}$, which could be derived through replacing B_{db}^{L} in Eq. (12) by B_{sb}^{L}) have been well bounded by the constraints from $B \to \pi K^{(*)}$ and ρK decays [19]. After considering the constraints from $B_s - \bar{B}_s$ mixing, which performs constraint on $b-s-Z'$ coupling solely, one may easily extract the results of $u-u-Z'$ and $d-d-Z'$ couplings $P_{ud}^{L,R}$ and $D_{ud}^{L,R}$ [21], which are recapitulated in Table 4[1]. Such results are the same as the ones in $b \to d\bar{q}q$ transition and will be adopted in the following analysis.

As for $b - d - Z'$ coupling, in Ref. [21], we have found the strength of B^L_{db} is strongly suppressed by $B_d - \bar{B}_d$ mixing. So, in our following fitting, the constraints from $B_d - \bar{B}_d$ mixing are also considered. The basic theoretical framework for $B_q - \bar{B}_q$ mixing within a family non-universal Z' model have been given by Refs. [20, 21] in detail. In this paper, we take the same framework and conventions as Ref. [21]. Moreover, we also take the recent updated fitting results of two mixing parameters C_{B_d} and ϕ_{B_d} by UTfit collaboration [23], which are defined as

$$C_{B_d} e^{2i\phi_{B_d}} \equiv \frac{\langle B_d|\mathcal{H}^{full}_{eff}|\bar{B}_d\rangle}{\langle B_d|\mathcal{H}^{SM}_{eff}|\bar{B}_d\rangle} = \frac{A^{SM}_d e^{i\phi^{SM}_d} + A^{NP}_d e^{i(2\phi^{NP}_d + \phi^{SM}_d)}}{A^{SM}_d e^{i\phi^{SM}_d}}\,, \tag{13}$$

and

$$C_{B_d} = 0.95 \pm 0.14\,, \quad \phi_{B_d} = -3.1^\circ \pm 1.7^\circ \tag{14}$$

at 68% probability[23].

3.2 Numerical analyses and discussions

With these theoretical formulae and the theoretical input parameters summarized in Appendix, we now present our numerical analyses and discussions. Our fitting is performed with that the experimental data is allowed within their 1.68σ ($\simeq$ 90% C.L.) error bars, while the theoretical uncertainties are also considered by varying the input parameters within their respective regions specified in the Appendix. In addition, we quote the scheme I to regulate the appearing end-point divergences. Since the Z' parameters $P^{L,R}_{ud}$ and $D^{L,R}_{ud}$ have been severely constrained by measured $B_s - \bar{B}_s$ mixing and $B \to \pi K$ decays [21], we demand that they cannot exceed their respective ranges listed in Table 4, where S1 and S2 correspond to UTfit collaboration's two fitting results for C_{B_s} and ϕ_{B_s} [23].

With $|B^L_{db}| = 0.15 \times 10^{-3}$ and the central values of theoretical input parameters, Fig. 2 shows the dependence of $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ on ϕ^L_d, $P^{L,R}_{ud}$ and $D^{L,R}_{ud}$. We find that $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ is

---

[1] In ref. [21], the errors for the numerical results of the Z' couplings are obtained simply with a Gaussian distribution treatment of the outputs caused by the uncertainties of input parameters. In this paper, to evaluate the conservative effects of the Z' couplings, we would study the whole possible regions of the Z' couplings, which are listed in Table 4.



Figure 2: The dependence of $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ on ϕ_d^L, $P_{ud}^{L,R}$ and $D_{ud}^{L,R}$ with $|B_{db}^L| = 0.15(\times 10^{-3})$ and the central values of theoretical input parameters.

Table 5: The numerical results for the Z' couplings.

Solutions	$\lvert B_{db}^L\rvert(\times 10^{-3})$	$\phi_d^L(°)$	P_{ud}^L	P_{ud}^R	D_{ud}^L	D_{ud}^R
S1	0.16 ± 0.08	-33 ± 45	0.45 ± 1.65	-1.2 ± 0.4	-0.70 ± 0.20	-0.02 ± 0.05
S2	0.19 ± 0.05	-50 ± 20	-0.31 ± 0.87	-0.68 ± 0.22	-0.41 ± 0.09	-0.02 ± 0.02

sensitive to the Z' contributions induced by P_{ud}^R and D_{ud}^L, but dull to P_{ud}^L and D_{ud}^R. Interestingly, as Fig. 2(b) and (c) shown, $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ could be reduced by Z' contributions with negative P_{ud}^R and D_{ud}^L at $\phi_d^L \sim -50°$, which possibly presents a solution for the large $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ problem.

With $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$, $A_{CP}^{dir}(\bar{B}_s \to \pi^- K^+)$, C_{B_d} and ϕ_{B_d} as constraints, corresponding to the two solutions for $P_{ud}^{L,R}$ and $D_{ud}^{L,R}$, the allowed regions for $|B_{db}^L|$ and ϕ_d^L are shown in Fig. 3



(a) (b)

Figure 3: The allowed regions for the parameters $|B_{db}^L|$ and ϕ_d^L in S1 (a) and S2 (b) under the constraints from $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ (blue region), $A_{CP}^{dir}(\bar{B}_s \to \pi^- K^+)$ (green region), C_{B_d} (brown region), ϕ_{B_d} (yellow region) and their combination (orange region) respectively.

and the numerical results are listed in Table 5. Interestingly, we find that the new fitting result $\phi_d^L = -33^\circ \pm 45^\circ$ ($-50^\circ \pm 20^\circ$) agrees well with the recent evaluation $\phi_d^L \sim -48^\circ$ [31] through the constraints from $B \to \pi\pi$. As found in Refs. [20, 21], $B_d - \bar{B}_d$ mixing performs a very strong constraint on B_{db}^L, whose magnitude is suppressed to be $< 0.24 \times 10^{-3}$ in both S1 and S2. While, the lower bound $0.08(0.14) \times 10^{-3}$ in S1(S2) for $|B_{db}^L|$ is dominated by $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$. Unfortunately, the constraint from $A_{CP}^{dir}(\bar{B}_s \to \pi^- K^+)$ is weak due to its large experimental error-bar. As shown in Table 4, since $P_{ud}^{L,R}$ in S2 are restricted to be smaller than their values in S1 case[21], the lower bound for $|B_{db}^L|$ in S2 is demanded larger than the one in S1. As shown in Fig. 3(a) and (b), the allowed ranges for both $|B_{db}^L|$ and ϕ_d^L in S2 are much smaller than the ones in S1. So, the solution S2 is much easier to be excluded by the future more precise measurements. Due to the constraint from $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$, the flavor-conserving Z' couplings $P_{ud}^{L,R}$ and $D_{ud}^{L,R}$ are further restricted. Their numerical results are also listed in Table 5.

With the values of Z' couplings listed in Table 5 and the other inputs parameters given in the Appendix, including their uncertainties, we present our theoretical predictions for the observables in the fifth and sixth column of Table 1, 2 and 3. We find $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ is reduced by Z' contributions. Due to large theoretical uncertainty, its lower bound 5.8(6.3)

$\times 10^{-6}$ agrees with experimental measurement $(5.0 \pm 1.1) \times 10^{-6}$ at 1σ error level in S1(S2). As a favored solution, within the allowed range for Z' couplings, we take

$$|B^L_{db}| = 0.24 \times 10^{-3}\,, \quad \phi^L_d = -55^\circ\,, \quad P^R_{ud} = -1.6\,(-0.9)\,, \quad D^L_{ud} = -0.90\,(-0.50)\,, \tag{15}$$

in S1(S2). With these Z' values and the central values of the other parameters as inputs, we get

$$\mathcal{B}(\bar{B}_s \to \pi^- K^+) = 7.8\,(8.5) \times 10^{-6}\,. \tag{16}$$

Comparing with the central value of the SM prediction 9.5×10^{-6}, we find $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ could be reduced by about 18% (11%) by Z' contributions at most. Such results means that S1 is more favored for the large $\mathcal{B}^{th}(\bar{B}_s \to \pi^- K^+)$ problem than S2. However, such result $\mathcal{B}(\bar{B}_s \to \pi^- K^+) = 7.8 \times 10^{-6}$ in S1 is also 2σ larger than the experimental result $(5.0 \pm 1.1) \times 10^{-6}$.

Our evaluations are based on the updated UTfit fitting results [23] for $B_q - \bar{B}_q$ mixing. Due to the constraint from $B_q - \bar{B}_q$ mixing, a CKM-like hierarchy is demanded in family non-universal Z' couplings, *i.e.*, $|B_{db}/B_{sb}| \sim |V^*_{td}V_{tb}/V^*_{ts}V_{tb}| \sim 0.2$ [21]. It is noted that both CDF and D0 collaborations have updated their measurements of $B_s - \bar{B}_s$ mixing recently based on $5.2 fb^{-1}$ and $6.1 fb^{-1}$ integrated luminosity, respectively. Their results read

$$\beta_s = [0.02, 0.52] \cup [1.08, 1.55] \quad 68\%\text{C.L.} \quad \text{CDF}\,[32]\,, \tag{17}$$

$$\phi_s^{J/\psi\phi} = -0.76^{+0.38}_{-0.36} \pm 0.02 \quad 68\%\text{C.L.} \quad \text{D0}\,[33]\,. \tag{18}$$

For comparison, D0 result Eq. (18) could also be written as

$$\beta_s = 0.38^{+0.18}_{-0.19} \pm 0.01 \quad 68\%\text{C.L.} \quad \text{D0}\,. \tag{19}$$

Unfortunately, these results have not been combined together yet by the two collaborations and by groups like UTfit [23], CKMfitter [34, 35], or HFAG [36]. It is interesting to note that, different from their former combined result $[0.27, 0.59] \cup [0.97, 1.30]$ [37], SM expectation for $\beta_s \sim 0.018$ agrees with CDF measurement Eq. (17) at $\sim 1\sigma$ level. D0 result in Eq. (19), which is similar to their former result $[0.27, 0.59] \cup [0.97, 1.30]$ [37], is also larger than SM prediction for β_s at $\sim 2\sigma$ level. So, obviously, CDF and D0 measurements have significantly difference with each other at small β_s region, where is important for constraining the strength of the NP contributions.



Figure 4: The allowed regions for the parameters $|B^L_{sb}|$ and ϕ^L_s under the constraints $\beta_s = [0.02, 0.52](CDF)$ (green region), ΔM_s (pink region) and their combination (blue region). The dashed line corresponds to the assumption $|B^L_{sb}| \approx |B^L_{db}| \approx 0.16 \times 10^{-3}$ as a special scenario.

If $\beta_s \sim 0.02$ is confirmed by the coming experimental measurement, CKM-like hierarchy in Z' couplings matrix possibly could be violated, which implies that $|B_{sb}|$ could be as small as $|B_{db}|$. To confirm such a guess, we revisit $B_s - \bar{B}_s$ mixing within the family non-universal Z' model following the same track as Ref. [21]. With $\beta_s = [0.02, 0.52]$(CDF Eq. (17)) and $\Delta M_s = 17.77 \pm 0.12$ [36] as constraints and all of the central values of the parameters listed in appendix as input, the allowed regions for the new parameters $|B^L_{sb}|$ and ϕ^L_s are shown by Fig. 4. So, from Fig. 4, one may easily find that the relation $|B_{sb}| \sim |B_{db}| \sim 0.2 \times 10^{-3}$, which is smaller than the former result $\sim 1 \times 10^{-3}$ [21] by a factor about 1/5, is allowed by the CDF updated measurement on β_s. Due to the constraints from $B \to \pi K$, ρK and πK^*, the values of the combining parameters $\zeta^{LL,LR}_s$ and $\xi^{LL,LR}_s$, whose form could be derived through replacing B^L_{db} in Eq. (12) by B^L_{sb}, are unchangeable [19, 21]. So, with $|B_{sb}| \sim |B_{db}| \sim 0.2 \times 10^{-3}$ as input, after reevaluating the constraints from $B \to \pi K$ *et al.* (as Refs. [19, 21] do), one may easily find that the results for $|P^{L,R}_{ud}|$ and $|D^{L,R}_{ud}|$ could be larger than the ones in Table 4 by a factor 5. So, as a possible special scenario, we take

$$|B^L_{db}| = 0.16 \times 10^{-3}\,, \quad \phi^L_d = -55^\circ\,, \quad P^R_{ud} = -1.2 \times 5\,, \quad D^L_{ud} = -0.70 \times 5\,, \tag{20}$$

for S1. With such Z' coupling values and the central values of other input parameters, we get $\mathcal{B}(\bar{B}_s \to \pi^- K^+) = 5.1 \times 10^{-6}$ which is perfectly consistent with the experimental data

$(5.0 \pm 1.1) \times 10^{-6}$. Interestingly, as found in Ref. [31], $|B_{sb}| \sim |B_{db}|$ is also needed for resolving the "$\pi\pi$ puzzle". It also should be noted that our above analysis is based on the data $\beta_s \sim 0.02$ is allowed, which is also needed to be confirmed by the future refined measurements of $B_s - \bar{B}_s$ mixing system at LHC-b and Tevatron.

4 Conclusion

In this paper, we have revisited $\bar{B}_s \to \pi K$, πK^* and ρK decays within QCDF formalism. We find that the large $\mathcal{B}^{th}(\bar{B}_s \to \pi^- K^+)$ problem could not be resolved by tuning the annihilation contributions. In order to pursue possible solutions for the large $\mathcal{B}^{th}(\bar{B}_s \to \pi^- K^+)$ problem, we evaluate the effects of a family non-universal Z' boson. $B_d - \bar{B}_d$ mixing is used to constrain the $b - d - Z'$ coupling. Our main conclusions are summarized as:

- The magnitude for $b - d - Z'$ coupling B_{db}^L is suppressed to $\sim 0.2 \times 10^{-3}$ by $B_d - \bar{B}_d$ mixing. Based on the constrained $u - u - Z'$ and $d - d - Z'$ couplings by $B_s - \bar{B}_s$ mixing and $B \to \pi K$ decays in Ref. [21], we find $P_{ud}^{L,R}$ and $D_{ud}^{L,R}$ are further restricted by $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$, especially P_{ud}^R and D_{ud}^L.

- A new weak phase $\phi_d^L \sim -50^\circ$, negative P_{ud}^R and/or D_{ud}^L with larger absolute value are helpful to improve the agreement of $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ between the SM prediction and the experimental measurement. Compared the SM result, $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ could be reduced by about 18%(11%) at most in S1 (S2). S1 is more favored by the large $\mathcal{B}^{th}(\bar{B}_s \to \pi^- K^+)$ problem than S2.

- The experimental measurement of β_s is very important to fix whether a CKM-like hierarchy is held in a family non-universal Z' couplings matrix. If the lower bound 0.02 for β_s measured by CDF is confirmed, $|B_{sb}|$ would be as small as $|B_{db}|$. Within such a scenario, $\mathcal{B}(\bar{B}_s \to \pi^- K^+)$ could be reduced to 5.1×10^{-6} which agrees with the experimental data $(5.0 \pm 1.1) \times 10^{-6}$.

The refined measurements for the B_s non-leptonic decay (especially $\bar{B}_s \to \pi^- K^+$) and $B_s - \bar{B}_s$ mixing in the LHC-b will provide a powerful testing ground for the SM and possible NP scenarios. Our analysis about the Z' effects on the observables for $\bar{B}_s \to \pi K$, πK^*, ρK

decays and $B_q-\bar{B}_q$ mixing are useful for probing or refuting the effects of a family non-universal Z' boson.

Acknowledgments

The work is supported by the National Science Foundation under contract No.11075059 and the Startup Foundation for Doctors of Henan Normal University under contract No.1006.

Appendix: Theoretical input parameters

For the CKM matrix elements, we adopt the UTfit collaboration's fitting results [23, 38]

$$\begin{aligned}
\overline{\rho} &= 0.132 \pm 0.02\,(0.135 \pm 0.04), \quad \overline{\eta} = 0.367 \pm 0.013\,(0.374 \pm 0.026), \\
A &= 0.8095 \pm 0.0095\,(0.804 \pm 0.01), \quad \lambda = 0.22545 \pm 0.00065\,(0.22535 \pm 0.00065). \qquad (21)
\end{aligned}$$

The values given in the bracket are the CKM parameters with assumption of the presence of generic New Physics, which are used in our evaluations when the Z' contributions are included.

As for the quark masses, we take the current quark masses

$$\begin{aligned}
\frac{\overline{m}_s(\mu)}{\overline{m}_q(\mu)} &= 27.4 \pm 0.4 \;\; [39], \quad \overline{m}_s(2\,\mathrm{GeV}) = 87 \pm 6\,\mathrm{MeV} \;\; [39], \quad \overline{m}_c(\overline{m}_c) = 1.27^{+0.07}_{-0.11}\,\mathrm{GeV}\; [40] \\
\overline{m}_b(\overline{m}_b) &= 4.20^{+0.17}_{-0.07}\,\mathrm{GeV}\; [40]\,, \quad \overline{m}_t(\overline{m}_t) = 164.8 \pm 1.2\,\mathrm{GeV}\; [40]\,, \qquad (22)
\end{aligned}$$

where $\overline{m}_q(\mu) = (\overline{m}_u + \overline{m}_d)(\mu)/2$, and the difference between u and d quark is not distinguished.

The other one is the pole quark mass. In this paper, we take [40, 41]

$$\begin{aligned}
m_u = m_d = m_s = 0, &\quad m_c = 1.61^{+0.08}_{-0.12}\,\mathrm{GeV}, \\
m_b = 4.79^{+0.19}_{-0.08}\,\mathrm{GeV}, &\quad m_t = 172.4 \pm 1.22\,\mathrm{GeV}. \qquad (23)
\end{aligned}$$

and the decay constants

$$\begin{aligned}
f_{B_s} &= (231 \pm 15)\ \mathrm{MeV}\ \ [42]\,, f_\pi = (130.4 \pm 0.2)\ \mathrm{MeV}\ [40]\,, f_K = (155.5 \pm 0.8)\ \mathrm{MeV}\ [40]\,, \\
f_{K^*} &= (217 \pm 5)\ \mathrm{MeV}\ [43]\,, f^T_{K^*}(2.2GeV) = (156 \pm 10)\ \mathrm{MeV}\ [43]\,, \\
f_\rho &= (209 \pm 2)\ \mathrm{MeV}\ [43]\,, f^T_\rho(2.2GeV) = (147 \pm 10)\ \mathrm{MeV}\ [43].
\end{aligned}$$

As for the B-meson lifetimes, $\tau_{B_s} = 1.470\,\mathrm{ps}$ [40] is used. We take the heavy-to-light transition form factors

$$A_0^{\bar{B}_s \to K^*}(0) = 0.360 \pm 0.034 \;[43]\,, F_0^{\bar{B}_s \to K}(0) = 0.30^{+0.04}_{-0.03} \;[44]\,. \tag{24}$$

References

[1] T. Altonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **103** (2009) 031801, arXiv:0812.4271 [hep-ex].

[2] CDF Collaboration,"Measurement of branching fractions and direct CP asymmetries of $B_s^0 \to h^+ H'^-$ decay in $1fb^{-1}$" and the updated results on April 10, 2008 may be found on http://www-cdf.fnal.gov/physics/new/bottom/bottom.html [Note 8579v1].

[3] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda , Phys. Rev. Lett. **83** (1999) 1914 [hep-ph/9905312]; Nucl. Phys. B**591** (2000) 313 [hep-ph/0006142].

[4] Y. Y. Keum, H. N. Li and A. I. Sanda, Phys. Lett. B **504** (2001) 6 [hep-ph/0004004]; Phys. Rev. D **63** (2001) 054008 [hep-ph/0004173].

[5] C. W. Bauer, S. Fleming and M. Luke, Phys. Rev. D **63** (2001) 014006 [hep-ph/0005275]; C. W. Bauer, S. Fleming, D. Pirjol and I. W. Stewart, Phys. Rev. D **63** (2001) 114020 [hep-ph/0011336]; C. W. Bauer and I. W. Stewart, Phys. Lett. B **516** (2001) 134 [hep-ph/0107001]; C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D **65** (2001) 054022 [hep-ph/0109045].

[6] M. Beneke and M. Neubert, Nucl. Phys. B **675** (2003) 333 [hep-ph/0308039].

[7] A. Ali, G. Kramer, Y. Li, C. D. Lu, Y. L. Shen, W. Wang and Y. M. Wang, Phys. Rev. D **76** (2007) 074018 [hep-ph/0703162].

[8] A. R. Williamson, J. Zupan, Phys. Rev. D **74** (2006) 014003, Erratum-ibid. D **74** (2006) 03901 [hep-ph/0601214].

[9] C. W. Chiang, M. Gronau, J. L. Rosner, Phys. Lett. B **664** (2008) 169, arXiv:0803.3229 [hep-ph].

[10] G. H. Zhu, JHEP **1005** (2010) 063, arXiv:1002.4518 [hep-ph].

[11] H. Y. Cheng and C. K. Chua, Phys. Rev. D **80** (2009) 114026, arXiv:0910.5237 [hep-ph].

[12] Y. G. Xu, R. M. Wang and Y. D. Yang Phys. Rev. D **79** (2009) 095017, arXiv:0903.0256 [hep-ph]; R. M. Wang and Y. G. Xu, Phys. Rev. D **81** (2010) 055011.

[13] G. Buchalla, G. Burdman, C. T. Hill and D. Kominis Phys. Rev. D **53** (1996) 5185 [hep-ph/9510376]; G. Burdman, K. D. Lane and T. Rador, Phys. Lett. B **514** (2001) 41 [hep-ph/0012073]; A. Martin and K. Lane, Phys. Rev. D **71** (2005) 015011 [hep-ph/0404107].

[14] E. Nardi, Phys. Rev. D **48** (1993) 1240 [hep-ph/9209223]; J. Bernabeu, E. Nardi and D. Tommasini, Nucl. Phys. B **409** (1993) 69 [hep-ph/9306251]; V. D. Barger, M. S. Berger and R. J. Phillips, Phys. Rev. D **52** (1995) 1663 [hep-ph/9503204]; M. B. Popovic and E. H. Simmons, Phys. Rev. D **62** (2000) 035002 [hep-ph/0001302]; T. G. Rizzo Phys. Rev. D **59** (1999) 015020 [hep-ph/9806397].

[15] M. S. Carena, A. Daleo, B. A. Dobrescu and T. M. P. Tait, Phys. Rev. D **70** (2004) 093009 [hep-ph/0408098].

[16] T. G. Rizzo, hep-ph/0610104; arXiv:0808.1906 [hep-ph].

[17] P. Langacker and M. Plümacher, Phys. Rev. D **62** (2000) 013006 [hep-ph/0001204].

[18] V. Barger, C. W. Chiang, P. Langacker and H. S. Lee, Phys. Lett. B **598** (2004) 218 [hep-ph/0406126].

[19] Q. Chang, X. Q. Li and Y. D. Yang, JHEP **0905** (2009) 056, arXiv:0903.0275 [hep-ph].

[20] V. Barger, L. L. Everett, J. Jiang, P. Langacker, T. Liu and C. E. M. Wagner, JHEP **0912** (2009) 048, arXiv:0906.3745 [hep-ph]; Phys. Rev. D **80** (2009) 055008, arXiv:0902.4507 [hep-ph].

[21] Q. Chang, X. Q. Li and Y. D. Yang, JHEP **1002** (2010) 082, arXiv:0907.4408 [hep-ph].

[22] A. Khodjamirian, T. Mannel and M. Melcher, Phys. Rev. D 68 (2003) 114007 [hep-ph/0308297]; Phys. Rev. D70 (2004) 094002 [hep-ph/0407226].

[23] M. Bona *et al.*, arXiv:0906.0953 [hep-ph]; M. Bona *et al.* (UTfit Collaboration) arXiv:0803.0659 [hep-ph]; online update at: http://www.utfit.org/UTfit/Results.

[24] G. Buchalla, A. J. Buras, and M. E. Lautenbacher, Rev. Mod. Phys. **68** (1996) 1125 [hep-ph/9512380].

[25] N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531; M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49** (1973) 652.

[26] T. Feldmann and T. Hurth, JHEP 11 (2004) 037 [hep-ph/0408188].

[27] J. M. Cornwall, Phys. Rev. D **26** (1982) 1453; J. Papavassiliou and J. M. Cornwall, Phys. Rev. D **44** (1991) 1285.

[28] Q. Chang, X. Q. Li and Y. D. Yang, JHEP **0809** (2008) 038, arXiv:0807.4295 [hep-ph].

[29] C. M. Zanetti and A. A. Natale, arXiv:1007.5072 [hep-ph].

[30] K. Cheung, C. W. Chiang, N. G. Deshpande and J . Jiang, Phys. Lett. B **652** (2007) 285 [hep-ph/0604223]; J. H. Jeon, C. S. Kim, J. Lee and C. Yu Phys. Lett. B **636** (2006) 270 [hep-ph/0602156]; C. H. Chen and H. Hatanaka, Phys. Rev. D **73** (2006) 075003 [hep-ph/0602140]; R. Mohanta and A. K. Giri, arXiv:0812.1842 [hep-ph]; J. Hua, C.S. Kim and Ying Li, Phys. Lett. B **690** (2010) 508 [hep-ph/0602156], arXiv:1002.2532 [hep-ph].

[31] Q. Chang, X. Q. Li and Y. D. Yang, arXiv:1003.6051 [hep-ph].

[32] CDF collaboration, "Updated Measurement of CP Violating Phase $\beta_s^{J/\psi\phi}$", Public Note 10206, May 10, 2010.

[33] D0 collaboration, "Updated Measurement of the CP-Violating Phase $\phi_s^{J/\psi\phi}$ Using Flavor-tagged Decay $B_s^0 \to J/\psi\phi$ ", Note 6098-CONF, July 22, 2010.

[34] CKMfitter Group (J. Charles et al.), Eur. Phys. J. C **41** (2005) 1 [hep-ph/0406184], updated results and plots available at: http://ckmfitter.in2p3.fr.

[35] A. Lenza *et.al.* (CKMfitter Group), arXiv:1008.1593 [hep-ph].

[36] E. Barberio *et al.* (Heavy Flavor Averaging Group), arXiv:0808.1297 [hep-ex], and online update at: http://www.slac.stanford.edu/xorg/hfag.

[37] CDF/D0 $\Delta\Gamma_s$,β_s Combination Working Group, "Combination of D0 and CDF Results on $\Delta\Gamma_s$ and the CP-Violating Phase $\phi_s^{J/\psi\phi}$", CDF Public Note 9787, D0 Note 5928-CONF, July 22, 2009.

[38] M. Bona *et al.* (UTfit Collaboration) JHEP **0507** (2005) 028 [hep-ph/0501199]; JHEP **0601** (2006) 081 [hep-ph/0606167]; online update at: http://www.utfit.org/UTfit/Results.

[39] Q. Mason *et al.* (HPQCD Collaboration), Phys. Rev. D **73** (2006) 114501 [hep-ph/0511160].

[40] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B **667**, 1 (2008).

[41] Tevatron Electroweak Working Group, CDF Collaboration and D0 Collaboration, arXiv:0808.1089 [hep-ex].

[42] Elvira Gamiz *et al.* (HPQCD Collaboration), Phys. Rev. D **80** (2009) 014503, arXiv:0902.1815 [hep-lat].

[43] P. Ball and R. Zwicky, Phys. Rev. D **71** (2005) 014015 [hep-ph/0406232]; **71** (2005) 014029 [hep-ph/0412079]; Phys. Lett. B **633** (2006) 289 [hep-ph/0510338].

[44] G. Duplancic and B. Melic, Phys. Rev. D **78** (2008) 054015, arXiv:0805.4170 [hep-ph].